Contents lists available at ScienceDirect

Tectonophysics

journal homepage: www.elsevier.com/locate/tecto

2019 M6.7 Yamagata-Oki earthquake in the stress shadow of 2011 Tohoku-Oki earthquake: Was it caused by the reduction in fault strength?



TECTONOPHYSICS

Keisuke Yoshida*, Naoki Uchida, Satoshi Hiarahara, Takashi Nakayama, Toru Matsuzawa, Tomomi Okada, Yoshiaki Matsumoto, Akira Hasegawa

Research Center for Prediction of Earthquakes and Volcanic Eruptions, Graduate School of Science, Tohoku University, Sendai, Japan

ARTICLE INFO ABSTRACT Earthquake occurrence in the stress shadow provides a unique opportunity for extracting information about the Keywords: Aseismic process physical processes behind earthquakes because it highlights processes other than the ambient stress change in Fluid migration earthquake generation. In this study, we examined the fault structure and the spatiotemporal distribution of the Stress shadow aftershocks of the 2019 M6.7 Yamagata-Oki earthquake, which occurred in the stress shadow of the 2011 M9.0 Hypocenter relocation Tohoku-Oki earthquake, to better understand the earthquake generation mechanism. The detection and hypo-Aftershock center relocation of the aftershocks led to the delineation of three planar earthquake structures consitent with 2016 Yamagata-Oki earthquake their focal mechanisms. The results suggest that individual aftershocks were caused by a slip on these macroscopic planar structures. Aftershock hypocenters rapidly migrated upward from the deeper part of the major plane (fault), similar to the recent earthquake swarm sequences following the Tohoku-Oki earthquake in the upper plate. Moreover, we investigated the temporal evolution of the surface strain rate distribution in the source region using GNSS data. The east-west contraction strain rate in the source region of the Yamagata-Oki earthquake, with an E-W compressional reverse fault mechanism, changed to an E-W extension as a result of the Tohoku-Oki earthquake, and it continued until the occurrence of the Yamagata-Oki earthquake. The upward hypocenter migrations, together with the earthquake occurrence in the stress shadow and in the E-W extension strain rate field, suggest that a reduction in the fault strength due to uprising fluids contributed to the occurrence of this earthquake sequence. Localized aseismic deformations, such as aseismic creeps, beneath the fault zone

may also have contributed to the earthquake occurrence. The results support the hypothesis that aseismic processes in the deeper part of the fault play crucial roles in the occurrence of shallow intraplate earthquakes.

1. Introduction

Several destructive earthquakes have occurred during the last 60 years in the active deformation zone along the eastern margin of the Japan Sea, such as the 1964 M7.5 Niigata earthquake, the 1983 M7.7 Nihonkai-Chubu earthquake, and the 1993 M7.8 Hokkaido Nansei-Oki earthquake (Fig. 1). Nakamura (1983) and Kobayashi (1983) hypothesized that this north-south striking deformation zone acts as a nascent plate boundary and releases the accumulated strain energy between the Eurasian and the North American plates. Subsequent seismological and space geodetic observations confirmed a relative plate motion proceeding along this deformation zone (Wei and Seno, 1998; Heki et al., 1999), although there are disagreements about the location and mode of this nascent boundary (Ohtake, 1995) and the microplate structures of both sides (Wei and Seno, 1998; Heki et al.,

1999). Complicated fault structures in this deformation zone suggest that the accumulated strain is released by multiple intraplate faults (Ohtake, 1995).

An M6.7 earthquake occurred west off the border between Yamagata and Niigata prefectures on June 18, 2019 (JST) in this deformation zone, referred to as the 2019 Yamagata-Oki earthquake (Figs. 1–3). The occurrence of this earthquake is enigmatic from the point of view of the temporal evolution of the stress field in this region that was affected by the 2011 M9 Tohoku-Oki earthquake. The state of stress in this deformation zone is the WNW–ESE compressional reversefaulting regime (Okamura et al., 1995). The 2011 Tohoku-Oki earthquake and the associated postseismic slip reduced this WNW–ESE compressional stress, and the reduced amount of differential stress was estimated to be \sim 0.5 MPa (Yoshida et al., 2012). Thus, the Tohoku-Oki earthquake created a stress shadow for typical WNW–ESE

E-mail address: keisuke.yoshida.d7@tohoku.ac.jp (K. Yoshida).

https://doi.org/10.1016/j.tecto.2020.228609

Received 23 June 2020; Received in revised form 24 August 2020; Accepted 25 August 2020 Available online 28 August 2020

0040-1951/ © 2020 Elsevier B.V. All rights reserved.



^{*} Corresponding author at: Research Center for Prediction of Earthquakes and Volcanic Eruptions, Tohoku University, 6-6 Aza-Aoba, Aramaki, Aoba-ku, Sendai 980-8578, Japan.



Fig. 1. Maps showing the source region of the 2019 M6.7 Yamagata-Oki earthquake. The red star indicates the hypocenter of the Yamagata-Oki earthquake. Black stars indicate the locations of the 1964 M7.5 Niigata earthquake, 1983 M7.7 Nihonkai-Chubu earthquake and its largest aftershock (M7.1), and 1993 M7.8 Hokkaido Nansei-Oki earthquake. Numbers above the stars denote the occurrence years and the JMA magnitudes. Orange lines indicate the traces of the active fault. Black thick lines and arrows indicate the plate boundaries and relative plate motions, respectively, according to Bird (2003). In this model, the location of the plate boundary between the Eurasian (or Amur) and North American (or Okhotsk) plates is based on Nakamura (1983). The broken curve indicates the location of this plate boundary proposed by Ohtake (1995). (a) Map showing northeast Japan. Blue arrows show the minimum compressional principle stress axis of the static stress change of the 2011 Tohoku-Oki earthquake (Yoshida et al., 2012). NA: North American plate, EU: Eurasian plate. Broken contour lines show the coseismic slip distribution of the 2011 Tohoku-Oki earthquake (Iinuma et al., 2012). (b) Enlarged view of the rectangle region of (a). Beach balls show the focal mechanisms listed in the F-net catalogue (Fukuyama et al., 1998). Gray and red dots show the hypocenters of shallow earthquakes (z < 40 km) with the JMA magnitude ≥ 2.0 before and after the 2011 Tohoku-Oki earthquake, respectively, for the period from Jan. 1, 1997 to August 20, 2019. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

compressional reverse-fault earthquakes in this deformation zone (Yoshida et al., 2019a). The shear stress magnitude on the fault of the Yamagata-Oki earthquake after the Tohoku-Oki earthquake should be lower than that just before the Tohoku-Oki earthquake. Nevertheless, the M6.7 reverse-fault earthquake occurred in the stress shadow after eight years (3021 d) from the 2011 earthquake (Fig. 3).

An important argument on the occurrence of shallow intraplate earthquakes is the existence of local sources affecting the states of stress and frictional strength of the fault zone (lio and Kobayashi, 2002; Calais et al., 2016). Previous studies suggested the important role of aseismic processes progressing in the deeper part of the fault, including pore pressure migrations and aseismic slips (Rice, 1992; Sibson, 1992; Liu and Zoback, 1997; Iio and Kobayashi, 2002; Hasegawa et al., 2005). Sibson (1992) proposed a model in which the pore pressure cycle associated with the upward fluid migrations controls the earthquake cycle of shallow intraplate earthquakes. In this fault-valve model, a local reduction in the fault strength due to increasing pore pressure is crucial for the occurrence of earthquakes. Iio and Kobayashi (2002) argued that the local stress concentration due to the localized deformation beneath the fault accounts for various geophysical and geological observations. Hasegawa et al. (2005) incorporated these ideas to the deformation model of the arc crust in the subduction zone, in which fluids dehydrated from the subducting slab play central roles.

The occurrence of the M6.7 earthquake in the stress shadow provides a unique opportunity for extracting information about the physical processes behind earthquake generation. This is because the occurrence of earthquakes in the stress shadow highlights the processes other than the ambient stress change in earthquake generation. Previous studies have analyzed several earthquake swarm sequences triggered in the stress shadow of the 2011 Tohoku-Oki earthquake (Fig. 1) and showed the upward migration behavior of hypocenters (Okada et al., 2015; Yoshida and Hasegawa, 2018a; Yoshida and Hasegawa, 2018b) along several planar structures; they hypothesized that swarm activities were triggered by the reduction in the fault strength due to the upward migration of the pore pressure despite the reduction in the shear stress. This hypothesis is consistent with the model proposed for northeast Japan, in which the slab-derived fluids that have reached the crust play a central role (Hasegawa et al., 2005).

In this study, we investigated the aftershock activities of the 2019 M6.7 Yamagata-Oki earthquake that occurred in the stress shadow of the 2011 M9.0 Tohoku-Oki earthquake to understand the earthquake generation mechanism. We detected early aftershocks, relocated the hypocenters and determined the focal mechanisms of earthquakes in the source region (Section 2). Moreover, we examined the temporal evolution of strain rates after the 2011 Tohoku-Oki earthquake. Afterwards, we investigated the fault structure and spatiotemporal evolution of the aftershocks (Section 3) and extracted information about the generation mechanism of the Yamagata-Oki earthquake and more generally, shallow intraplate earthquakes (Section 4).

2. Data and methods

In this section, we describe the data (Subsection 2.1) and methods to detect early aftershocks (Subsection 2.2), to relocate hypocenters



Fig. 2. Distribution of the hypocenters in the Japan Meteorological Agency (JMA) unified catalogue from March 1, 2003 to May 10, 2020. Blue circles represent the hypocenters. The size of each circle corresponds to the diameter of the fault, assuming a stress drop of 3 MPa. The star denotes the hypocenter of the mainshock. (a) A map view showing the focal region of the 2019 Yamagata-Oki earthquake and the hypocenter distribution. (b) A map view showing the focal mechanisms determined by the JMA by red "beach balls." (c)–(j) Cross-sectional views of vertical hypocenter distribution along the lines indicated in (a). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

(Subsection 2.3), and to determine focal mechanisms (Subsection 2.4) of earthquakes that occurred in the source region of the 2019 Yamagata-Oki earthquake. We also explain the method to estimate the surface strain rate prior to the mainshock by using global navigation satellite system (GNSS) data (Subsection 2.5).

2.1. Data

We analyzed 4255 earthquakes listed in the Japan Meteorological Agency (JMA) unified catalogue from March 1, 2003 to May 10, 2020 (Fig. 2) and the waveforms recorded at the seismic stations surrounding the source region (Fig. S1). For the following waveform analyses (Subsections 2.2, 2.3 and 2.4), we constructed P- and S-wave windows starting from 0.3 s before the arrival with the durations of 2.5 s and 4.0 s, respectively, for each earthquake-station pair. For the arrival time, we used manually picked times listed in the JMA unified catalogue, if available; otherwise, we used theoretical arrival times computed by assuming the same velocity model as used in the JMA unified catalogue. Moreover, we used the daily location data at the national GNSS network called GEONET (GNSS Earth Observation Network System; https://www.gsi.go.jp/ENGLISH/index.html) (Fig. S1) provided by the Geospatial Information Authority of Japan (GSI) to estimate the strain rate (Subsection 2.5).

2.2. Detection of early aftershocks

The minimum magnitude of catalogue completeness temporally increases for a certain period immediately after a large earthquake. To obtain a comprehensive view of the spatiotemporal evolution of the aftershocks of the 2019 Yamagata-Oki earthquake, we performed early aftershock detection to identify the undetected earthquakes in the JMA unified catalogue. In the first step, we used the waveform data obtained at the two nearest stations to identify the possible events, which are not detected in the JMA unified catalogue. In the second step, we used the waveform data obtained at a large number of stations to verify that the identified events are earthquakes in the source region.

We first applied the template matching method (Gibbons and Ringdal, 2006; Schaff and Waldhauser, 2010) to the continuous waveform data obtained at the two nearest stations (YATSUM and AWASHI shown by purple inverted triangles in Fig. S1) from the source region from June 18 to June 19, 2019 (JST). We followed the procedure outlined in Yoshida and Hasegawa (2018a) modified after Shelly et al. (2013). We used 4255 earthquakes in the JMA unified catalogue (referred to as JMA events) as template events. We applied a bandpass filter with a passband frequency of 5–12 Hz to the continuous waveforms and the S-wave windows of the template events and then downsampled the waveforms to 25 samples per second.

We compared the S-wave windows of the template events (duration of 4.0 s) with the continuous waveform data; we calculated the correlation coefficients cc_i (the *i*-th point) on each waveform at each time lag



Fig. 3. Magnitude-time diagrams of events in and around the source region of the 2019 Yamagata-Oki earthquake for (a) -50 d to 350 d. (b) -6000 d to 350 d. and (c) -1 dto 2 d following the mainshock. The blue and red circles with gray bars indicate the earthquake magnitudes listed in the JMA unified catalogue and those of newly detected in this study, respectively. The black line denotes the cumulative number of earthquakes with magnitudes greater than 2.0. The green vertical line indicates the occurrence time of the 2011 Tohoku-Oki earthquake. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

(incremented at 0.04 s intervals), We computed the time series of $cc_i' = \max(cc_{i-1}, cc_i, cc_{i+1})$, and obtained the sum of cc_i' over all the stations and components (CC_i). We computed the median value and the median absolute deviation (MAD) of CC_i for the day. If CC_i exceeds the median value added by the twelve times of MAD, we regarded it as a possible missing event unless it is listed in the JMA unified catalogue. Consequently, we newly obtained 1207 possible events.

As the second step, we eliminated uncertain events from the newlydetected possible events by comparing the waveforms of the events with those of the JMA events at various stations (46 green triangles in Fig. S1). For each possible event, we constructed the P-wave and Swave windows in the same way as for the JMA events (durations of 2.5 s for P-wave and 4.0 s for S-wave starting at 0.3 s before their onsets). The onset timing is assumed to be the same as that of the most correlated JMA event during the detection. We evaluated the waveform similarity between each newly-detected event and JMA event at each station and component; when a newly-detected event did not satisfy the waveform similarity threshold for five or more JMA events, we eliminated that event. The threshold of the waveform similarity is 0.8 in cross-correlation coefficients for the P- and/or S-wave windows at more than two stations. Consequently, we eliminated 829 events and obtained 378 new events (Fig. 3c). Thus, we created our earthquake catalogue, which consists of 4556 earthquakes in total.

We determined the magnitudes of the newly-detected events based on the relative waveform amplitudes of P- and S-waves to those of JMA events. We assumed that the JMA magnitude is equal to the moment magnitude of Hanks and Kanamori, 1979 and the ratio of the waveform amplitude is equal to the ratio of the seismic moment when the station is the same and the sources have similar focal mechanisms and locations. We first aligned the timings of the waveforms by using the lag time of each waveform pair with high similarity (cc ≥ 0.8). The waveform similarity ensures similarity in the source mechanism and location. Afterwards, we compared the amplitudes of the waveform pairs at each time, applied principal component analysis (PCA), and obtained the best amplitude ratio of the waveform pair (Yoshida et al., 2019b). We obtained the amplitude ratios of a detected event with various JMA events at various seismic stations and determined the seismic moment of the detected event as the mean value of individual estimates weighted by the cross-correlation coefficient. We finally determined the magnitude of the detected events by using the relationship between the seismic moment and the moment magnitude (Hanks and Kanamori, 1979). We estimated the completeness of magnitude (M_c) by the modified maximum curvature method reported in Woessner and Wiemer (2005). Our earthquake detection successfully reduced M_c from 1.3 to 1.0 for the day following the Yamagata-Oki earthquake (Fig. S3).

2.3. Hypocenter relocation

We relocated the hypocenters of earthquakes from our earthquake catalogue obtained in Subsection 2.2 using the double-difference method (Waldhauser and Ellsworth, 2000) following the procedure outlined in Yoshida and Hasegawa (2018a). We first extracted the Pwave (101,611) and S-wave (131,760) differential arrival time data from the JMA unified catalogue of the event pairs, with distances less than 4.0 km. We also used the waveform data obtained at stations close to the source area (Fig. S1) for the waveform correlation measurements of differential arrival times. These stations are operated by Tohoku University, JMA, and NIED Hi-net (National Research Institute for Earth Science and Disaster Resilience, 2019a) where three-component velocity-type seismograms are recorded at a sampling rate of 100 Hz. We applied a bandpass filter with frequencies ranging between 5 and 12 Hz and computed the cross-correlation function for the waveforms of each event pair, with a horizontal distance less than 3.0 km. Differential arrival times derived from the cross-correlation function were used for the relocation if the cross-correlation coefficient was higher than 0.8. The number of the differential arrival time data for P- and S-waves, derived from the waveform cross-correlation delay measurements, was 650,500 and 810,036, respectively. The mean number of correlated neighbors (at more than two stations) of each event was 63 for JMA events and 49 for newly-detected events.

We applied the double-difference earthquake relocation method (Waldhauser and Ellsworth, 2000) to the differential arrival time data. We assumed the 1-D velocity model of Hasegawa et al. (1978), which was routinely used at Tohoku University to determine the hypocenter locations and focal mechanisms of the events in northeast Japan. Hypocenters were updated during 15 iterations. In the first five iterations, we weighted the catalogue data 10 times larger than the cross-correlation derived data to constrain the relative locations at a larger scale (> 1 km). In the latter 10 iterations, we weighted the cross-correlation derived data 50 times stronger than the catalogue data to delineate shorter-scale (< 1 km) structures. We evaluated the uncertainty in the relative hypocenter locations by recalculating the hypocenters 1000 times, based on bootstrap resampling of differential arrival time data.

2.4. Determination of focal mechanisms

To determine the focal mechanisms, we used the relative amplitudes of direct P- and S-waves corrected by those of a reference earthquake whose focal mechanism was already determined. We followed the procedure outlined in Yoshida et al. (2019b), which utilizes the amplitude ratios of P-, SH-, and SV-waves by assuming that the medium in the vicinity of the source is homogeneous and isotropic (Dahm, 1996). We restricted the distance between the target and reference events to less than 3 km to reduce the differences in the propagation and site effects. We used the waveform correlation between the target and reference earthquakes to reliably obtain the amplitude ratio data.

We adopted twelve focal mechanisms listed in the JMA unified catalogue (Fig. 2b) as reference focal mechanisms using the P-wave first-motion polarity data. We applied a bandpass filter with frequency of 2–5 Hz to both the P-wave and S-wave windows. The amplitude ratio

was computed at each seismic station using the same method described in Subsection 2.1, when the cross-correlation coefficient is greater than 0.75 between the waveforms of the target event and the reference event. If the amplitude ratio data were obtained from more than eight different seismic stations and 15 channels, we estimated the moment tensor components. We computed 2000 focal mechanisms for each target event based on the bootstrap resampling of the amplitude ratio data. The difference in the focal mechanisms from the best solution was measured by the 3-D rotation angle (Kagan, 1991). If the 90% confidence interval was larger than 30°, we discarded the result. If focal mechanism of one target event was obtained from different reference events, we adopted the one with the minimum confidence interval.

2.5. Estimation of strain rate prior to the mainshock

We estimated the temporal changes in the surface strain rate prior to the 2019 Yamagata-Oki earthquake by using the location data at the national GNSS network GEONET. We estimated the spatial distribution of surface strain rates in the following periods: (1) March 9, 2009 to March 10, 2011 (before the 2011 Tohoku-Oki earthquake), (2) March 12, 2011 to June 18, 2014 (after the coseismic change of the 2011 Tohoku-Oki earthquake), (3) June 19, 2014 to June 18, 2018, and (4) June 19, 2018 to June 18, 2019 (before the 2019 Yamagata-Oki earthquake).

We used the routine daily coordinate solutions (F3 solutions) provided by the GSI obtained at the GEONET stations (Fig. S1). We applied the method by Shen et al. (1996) to the mean displacement rates during these periods and obtained the spatial distribution of the strain rate at each 0.05° -spaced grid node.

3. Results

3.1. Fault Structure

We obtained the relocated hypocenters of 4501 events and determined the focal mechanisms for 269 events. Location data for 55 earthquakes were removed during the relocation procedure because their hypocenters were located above the ground surface (49 JMA events and six newly-detected events). We computed the differences between the maximum and minimum values in the 95% confidence interval of the hypocenter locations (Fig. S3) and obtained the medians as a measure of estimation error of the relative location: 0.0014° in longitude (0.12 km in distance), 0.0013° in latitude (0.14 km in distance), and 0.22 km in depth for the JMA events and 0.0021° in longitude (0.18 km in distance), 0.0017° in latitude (0.19 km in distance), and 0.33 km in depth for the newly-detected events.

The present hypocenter relocation succeeded in delineating a few planar structures of hypocenters (Fig. 4) from the scattered distribution of the initial hypocenters of the JMA unified catalogue (Fig. 2). Such drastic changes in the hypocenter distribution were also obtained for other earthquake clusters in northeast Japan based on a similar relocation method by using numerous and precise differential arrival time data (Yoshida and Hasegawa, 2018a; Yoshida and Hasegawa, 2018b; Yoshida et al., 2020). The largest plane of aftershock hypocenters is the one dipping eastward approximately 15 km long and 10 km wide (gray rectangle in Fig. 4a and gray lines in Fig. 4c–i), at the bottom of which the mainshock hypocenter is located. We also observed two west-dipping conjugate planes in the southern part (green rectangle and green lines) and the shallower part (black rectangle and red lines) of the aftershock area.

Some earthquakes appear to have occurred along the same fault structures as aftershocks even before the mainshock. A few of them were located near the hypocenter of the mainshock, but most were located on the western segments of the fault structure of aftershocks (Fig. 5c and d).

Fig. 6 shows the spatial distribution of the focal mechanisms



Fig. 4. Distribution of the relocated hypocenters. (a) A map view showing the focal region of the 2019 Yamagata-Oki earthquake. Blue circles represent the hypocenters. The size of each circle corresponds to the diameter of the fault, assuming a stress drop of 3 MPa. The star denotes the hypocenter of the mainshock. Gray, green, and red rectangles denote the model fault planes. Red crosses indicate the hypocenters of the newly-detected events. The sizes of the crosses are ten times as large as those of the JMA events in order to clearly show their distribution. (b) The moment release amount of the aftershocks computed at each 0.04° -spaced grid-cel is shown by the colour scale. The black ellipse shows the seismic gap. (c)–(j) Cross-sectional views of the vertical hypocenter distribution along the lines indicated in (a). Gray, green, and black lines denote the model fault planes. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

determined in this study and by the JMA before and after the Yamagata-Oki earthquake. Most of them have WNW–ESE compressional reversefault focal mechanisms similar to that of the mainshock. Most of those aftershocks have nodal planes nearly parallel to the alignments of the hypocenters, suggesting that individual aftershocks occur on the macroscopic planar structures illustrated by the hypocenters.

3.2. Seismic gap of earthquakes before and after the mainshock

Aftershocks did not occur in the area up to 2–3 km from the mainshock hypocenter along the east-dipping fault plane (z = 12.5-15.0 km in Fig. 4f–h). Intensive aftershock activities occurred in the shallower portions above this seismic gap along the east-dipping plane (z = 10.0-12.5 km) and along two conjugate planes. A seismic gap of aftershocks formed at a deeper portion of the aftershock area and along the shallower side of the mainshock hypocenter. In this area, earthquakes did not occur before the mainshock.

To analyze this tendency differently, we computed the seismic moment release amount of each earthquake by assuming that the magnitude is equal to the moment magnitude. Then, we summed the moment release amount at each evenly-spaced point every 0.04° by using the aftershocks within the nearest grid-cell (Fig. 4(b)). The moment release amount of aftershocks and pre-mainshock earthquakes is smaller ($< 10^{11} - 10^{12}$ Nm) in the region up to 2–3 km from the

main shock hypocenter along the east-dipping fault plane, corresponding to the seismic gap, than the other regions in its shallower extension ($\sim 10^{13}$ Nm).

3.3. Migration of the aftershock distribution

The aftershock area appears to migrate with time (Fig. 5c and d). Fig. 7 shows the spatiotemporal evolution of the aftershock distribution in four periods: (a), (b) 0 to 0.2 d, (c), (d) 0.2 to 2 d, (e), (f) 2 to 30 d, and (g), (h) 30 to 300 d from the mainshock. Hypocenters tend to be located in the deeper portion in the earlier period (Fig. 7a and b) but expand to the shallower part in the later periods (Fig. 7c through d). Fig. 8 shows the direct comparison of the occurrence times of aftershocks with depth, longitude, and latitude. We sorted the earthquakes by time and divided them into 45 bins with 100 events. We computed the two levels above or below where 10% of the events are contained in each bin and showed the levels in Fig. 8. The aftershock hypocenters moved toward the shallower part approximately following the logarithm of time (Fig. 8a and b). Aftershocks in the deeper portion of the fault (z > 13 km) intensively occurred in the first day from the mainshock but ceased after ~1 d from the mainshock (Fig. 8a).

Aftershock migration is clearly seen, especially for the period 0.3 days after the mainshock, during which M_c was relatively high (1.3), even after detection (Fig. S3c). One concern is that aftershock



Fig. 5. Spatiotemporal distribution of earthquakes. The occurrence times measured from the mainshock are shown in the colour scale in (a), (c) map-view and (b), (d) cross-sectional view within the range shown by the broken rectangle in (a) and (c). (a), (b): before the mainshock. (c), (d): after the mainshock. The star indicates the mainshock.

migration is actually an outcome of the temporal change in the detection capability. However, the same aftershock migration is observed even if we set the lower limit of magnitude to be 1.5 (Fig. S4), which is higher than M_c for the first day after the mainshock. Moreover, aftershock migration is visible not only in the first 0.3 days, but also in the 100 days following the mainshock (Fig. 8b). These observations support that the observed aftershock migration is not artifact by the temporal change in the detection capability, but actually occurred.

3.4. Strain rate prior to the mainshock

Fig. 9 shows the spatial distribution of the strain rate for four periods: (a) March 9, 2009 to March 10, 2011 (before the 2011 Tohoku-Oki earthquake), (b) March 12, 2011 (after the coseismic change of the 2011 Tohoku-Oki earthquake) to June 18, 2014, (c) June 19, 2014 to June 18, 2018, and (d) June 19, 2018 to June 18, 2019 (before the 2019 Yamagata-Oki earthquake).

Northeast Japan including the source region of the Yamagata-Oki earthquake was characterized by the E–W contraction strain rate before the 2011 Tohoku-Oki earthquake (Fig. 9a), which is consistent with the results of the previous studies (Sagiya et al., 2000; Yoshida et al., 2015). However, the 2011 Tohoku-Oki earthquake drastically changed the spatial distribution of the strain rate, and the east-west extension strain rate spread all over the region (Fig. 9b). This E–W extension continued near the source region of the Yamagata-Oki earthquake after the Tohoku-Oki earthquake (Fig. 9c) because of the postseismic deformation,

such as the afterslip along the plate interface (Ozawa et al., 2011) and the viscoelastic response of the mantle (Sun et al., 2014; Hu et al., 2016; Tomita et al., 2020). The E–W extension strain rate further continued until the occurrence of the Yamagata-Oki earthquake, as shown in the strain rate map for the latest period (from June 19, 2018 to June 18, 2019), which was just before the Yamagata-Oki earthquake.

There are two GNSS stations near the source region of the Yamagata-Oki earthquake (blue crosses in Fig. S1), one of which is located west of the source region and the other east. Fig. 10 shows the temporal changes in the distance in the E–W direction between these stations. The E–W distance between the two stations continued to increase after the 2011 Tohoku-Oki earthquake, suggesting that the E–W extension continued even before the Yamagata-Oki earthquake.

4. Discussion

4.1. Occurrence mechanism of the 2019 Yamagata-Oki earthquake in the stress shadow

The 2019 M6.7 Yamagata-Oki earthquake occurred in the coseismic stress shadow of the 2011 M9.0 Tohoku-Oki earthquake. The seismicity rate in the source region of the Yamagata-Oki earthquake decreased after the Tohoku-Oki earthquake in agreement with the stress shadow (Fig. 3b). The source region was located in the E–W extension and north–south contraction strain rate field even immediately before the Yamagata-Oki earthquake (Fig. 9d). The occurrence of the Yamagata-



Fig. 6. (a), (b) Distribution of the focal mechanisms of aftershocks. Focal mechanisms are shown by the beach ball in (a), (c) map-view and (b), (d) cross-sectional view within the range shown by the broken rectangle in (a) and (c). (a), (b): before the mainshock. (c), (d): after the mainshock. Blue circles indicate the locations of relocated hypocenters.

Oki earthquake with the east-west compressional reverse-faulting mechanism in this situation provides important constraints for the occurrence mechanism, on the basis that there need to be some local effects to cause an earthquake in the macroscopic stress shadow; thus, the occurrence of this earthquake is probably related to the local effects on the stress and/or fault strength. We observed an upward migration of aftershocks of the 2019 Yamagata-Oki earthquake (Fig. 8b). Similar upward migrations were also observed for several earthquake swarms triggered in the central part of northeast Japan after the 2011 Tohoku-Oki earthquake (Kosuga et al., 2014, Okada et al., 2015; Yoshida and Hasegawa, 2018a, Yoshida and Hasegawa, 2018b). Those earthquake swarms occurred in the stress shadow of the 2011 Tohoku-Oki earthquake as well and were caused by the increase in pore pressure after the 2011 earthquake (Terakawa et al., 2013; Yoshida et al., 2016). Many aftershocks of the Yamagata-Oki earthquake occurred on the three dominant planar structures, which may indicate that the planes are highly permeable and acted as fluid conduit. In the later period, some aftershocks are located above the three dominant planes (8 < z < 10 km in Figs. 7 and 8), which may have occurred on small fault structures activated by upward fluid discharge from the dominant planes. Although we could not estimate the focal mechanisms of the shallower earthquakes, some of the shallow events may have focal mechanisms unfavorably oriented from the regional stress field due to high pore pressures (Terakawa et al., 2013; Yoshida et al., 2016). By considering the involvement of fluids, we can also explain the occurrence of the mainshock in the stress shadow because the increase in

pore fluid pressure reduces the fault strength.

Aftershock migration can be explained by the effects other than the pore pressure migration, such as postseismic slip propagation (Wesson, 1987; Kato, 2004; Hsu et al., 2006; Kato, 2007; Peng and Zhao, 2009; Frank et al., 2017; Perfettini et al., 2018; Yoshida et al., 2020) and aftershock-to-aftershock interactions (Helmstetter and Sornette, 2002). By adopting these models, we may explain the observed upward aftershock migration without pore pressure migration. The expansion of the aftershock region with the logarithm of time observed in this study (Fig. 8b) is consistent with the observations and simulations of the postseismic slip (Kato, 2007; Peng and Zhao, 2009; Perfettini et al., 2018) and difficult to explain by the commonly used simple pore pressure diffusion model (Shapiro et al., 1997). However, if these stress triggering models are adopted as the cause of the migrationof the aftershock region, the mainshock occurrence in the stress shadow should still be explained independently. We also need to explain the occurrence of the M6.7 E-W compressional reverse-faulting earthquake in the field with the increased E-W extensional strain rate.

One possible explanation is based on the localized deformation in the deeper part; if aseismic slip proceeds in the deeper extension of the fault, they may locally increase the shear stress on the fault and cause an earthquake even in the macroscopic stress shadow of the Tohoku-Oki earthquake. Meneses-Gutierrez and Sagiya (2016) recently analyzed the surface deformation in the Niigata-Kobe Tectonic Zone (NKTZ), which is also located in the stress shadow of the 2011 Tohoku-Oki earthquake. They found a localized contraction zone around the



9

(caption on next page)

Fig. 7. Spatiotemporal distribution of earthquakes for four periods: (a), (b) 0 to 0.2 d, (c), (d) 0.2 to 2 d, (e), (f) 2 to 30 d, and (g), (h) 30 to 300 d from the mainshock. The colour scale in shows the relative number of earthquakes ordered by time from the beginning to the end of the periods. (a), (c), (e), (g): map-view; (b), (d), (f), (h): cross-sectional view within the range shown by the broken rectangle in (a), (c), (e) and (g). Gray and black circles show the location of earthquakes before and after these periods, respectively.

northern part of the NKTZ after applying a short wavelength filter both before and after the Tohoku-Oki earthquake and explained this observation by a persistent aseismic creep in the deeper portion of the fault. Aseismic creeps may decelerate when the shear stress magnitude decreases but may not completely cease. Previous studies suggested that the active deformation zone along the eastern margin of the Japan Sea, where the Yamagata-Oki earthquake occurred, is the northern extension of the NKTZ (Nakamura, 1983). Thus, aseismic creep might have also occurred in the deeper extension of the source fault of the Yamagata-Oki earthquake. However, it is not likely that such a deeper deformation alone increased the shear stress on the fault to exceed the level just before the 2011 Tohoku-Oki earthquake and caused the mainshock rupture of the 2019 Yamagata-Oki earthquake since such aseismic slips should have slowed down after the 2011 earthquake even if they existed. Furthermore, there is no observational evidence that such deep extension of the fault actually exists in the source area of the Yamagata-Oki earthquake. In fact, we did not observe an east-west contraction between the two nearest GNSS stations before the Yamagata-Oki earthquake (Fig. 10).

Thus, we preferred the pore pressure migration model to explain the occurrence of the 2019 Yamagata-Oki earthquake. However, if a deep extension of the fault actually existed, pore pressure increase may also promote localized deformation beneath the fault zone. It may reduce the strength of the deeper extension part of the fault, accelerate creeps on it and even trigger aseismic slip on it. It is possible that pore pressure migration and localized deformation in the deeper extension of the fault may have coexisted and contributed to the occurrence of the 2019 Yamagata-Oki earthquake. The propagation of the postseismic slip and the aftershock-to-aftershock interaction may also have contributed to the aftershock migration.

Seismic tomography results suggest that a low-velocity zone in the mantle wedge extends westward from beneath the volcanic front to beneath the Japan Sea (Zhao et al., 2011). The low-velocity zone is considered to represent the ascending flow portion of the secondary convection within the mantle wedge and, therefore, contains fluids dehydrated from the slab and resultant melts (Hasegawa and Nakajima, 2004). Fluids released from the mantle diapirs rising from this low-velocity zone in the mantle wedge may reach beneath the focal region of the 2019 Yamagata-Oki earthquake. They may also contribute to the formation of the active deformation zone along the eastern margin of the Japan Sea.

The present observations support the hypothesis that the involvement of fluids plays an important role in the occurrence of the mainshock-aftershock sequences. The involvement of fluids has been more often reported for earthquake swarms (Parotidis et al., 2005; Yukutake et al., 2011; Shelly et al., 2013) but has also been considered for mainshock-aftershock sequences (Nur and Booker, 1972; Yamashita and Knopoff, 1987; Sibson, 1992). In fact, the differential stress magnitude in northeast Japan is estimated to be too small (a few tens of MPa) to cause earthquakes without the reduction in fault strength (Yoshida et al., 2014; Yoshida et al., 2015). Recently, Matsumoto et al. (2020) examined the foreshocks and aftershocks of the 2017 M5.3 Kagoshima Bay earthquake and reached a similar conclusion that the foreshock-mainshock-aftershock sequences were caused by the fluid movements. They found that foreshock hypocenters migrated along the mainshock fault plane and that aftershock hypocenters migrated upward similarly to the present observations. In the fault-valve model of Sibson (1992), uprising fluids originated from the deeper portion reduced the fault strength, caused the earthquake, and discharged further upward. This model not only accounts for the observation features of the 2017 Kagoshima Bay earthquake, but also the newly observed features of the 2019 Yamagata-Oki earthquake; i.e., its occurrence in the stress shadow and the upward migration of aftershock hypocenters.

4.2. Seismic gap and the large coseismic slip region of the mainshock

We delineated three planar structures of hypocenters (Fig. 4). The mainshock hypocenter is located at the deepest edge of the east-dipping plane of the hypocenters distribution, which likely represents the mainshock fault plane, on which a seismic gap (the ellipse in Fig. 4b) existed near the mainshock hypocenter.

Previous studies reported a spatial separation between the mainshock coseismic slip area and the areas where aftershocks are active based on the direct comparison of aftershock distribution and coseismic slip distribution (Mendoza and Hartzell, 1988; Das and Henry, 2003; Woessner et al., 2006; Asano et al., 2011; Ebel and Chambers, 2016; Yoshida et al., 2020; Wetzler et al., 2018). The seismic gap of aftershocks near the mainshock hypocenter (the ellipse in Fig. 4b) likely represents a large coseismic slip area of the mainshock (Fig. 4a and b). If we assume that the aftershock gap represents the large slip region, the length (~11.5 km) is much smaller than the typical fault length of a M6.7 earthquake, based on the circular crack model (Eshelby, 1957) with a stress drop of 3 MPa (~25 km) and a model based on empirical relationships (~30 km; Wells and Coppersmith, 1994), suggesting that a substantial portion of seismic moment was released on the small asperity. However, the result of a waveform inversion based on strong motion data from NIED K-NET and Kik-net (National Research Institute for Earth Science and Disaster Resilience, 2019b) of the Yamagata-Oki earthquake indicates that the large coseismic slip of this earthquake actually occurred in the area with many aftershocks (https://www. kyoshin.bosai.go.jp/kyoshin/topics/Yamagataoki_20190618/

inversion_en/) west from the aftershock gap, although the discrepancy between the region with large coseismic slip and the aftershock gap comes from the estimation error of the coseismic slip distribution due to the difficulty in constraining the rupture speed (Lay et al., 2010).

5. Conclusions

In this study, we investigated the fault structure and spatiotemporal variations of the aftershocks of the 2019 M6.7 Yamagata-Oki earthquake, which occurred in the stress shadow of the 2011 M9.0 Tohoku-Oki earthquake, to understand the occurrence mechanisms of earthquake sequences. We exploited the assumption that there need to be some local effects to cause an earthquake in the macroscopic stress shadow. Additionally, we investigated the spatiotemporal evolution of horizontal strain rates before the occurrence of the mainshock.

Precisely relocated aftershock hypocenters formed three planar structures. One of the two nodal planes of focal mechanisms of each event were almost parallel to the hypocenter alignments, suggesting that individual aftershocks were mainly caused by afterslip along the faults delineated by the aftershocks. A seismic gap of aftershocks near the mainshock hypocenter on the major east-dipping plane may reflect the area of large coseismic slip during the mainshock. Aftershock hypocenters rapidly migrated upward from the deeper part of the fault, similar to the recent earthquake swarm sequences triggered after the 2011 Tohoku-Oki earthquake in the stress shadow.

The spatial distribution of the surface strain rate in northeast Japan, including the source region of the 2019 Yamagata-Oki earthquake, was characterized by an E–W contraction strain rate; however, the 2011 Tohoku-Oki earthquake drastically changed the spatial distribution and



Fig. 8. Spatiotemporal distribution of aftershocks. (a), (b): depth, (c), (d): longitude, and (e), (f): latitude. Blue and red circles show the events taken from the JMA unified catalogue and those detected in this study, respectively. Occurrence times are shown in the linear scale in (a), (c), and (e) and in the logarithmic scale in (b), (d), and (f). Yellow curves in (b), (c), and (d) show the levels above and below where 10% of the events are contained in each bin with 100 events sorted by time. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)



Fig. 9. Spatial distribution of the strain rate in northeast Japan obtained from GEONET data for four periods prior to the 2019 Yamagata-Oki earthquake: (a) March 9, 2009 to March 10, 2011 (before the 2011 Tohoku-Oki earthquake), (b) March 12, 2011 (after the coseismic change of the 2011 Tohoku-Oki earthquake) to June 18, 2014, (c) June 19, 2014 to June 18, 2018, and (d) June 19, 2018 to June 18, 2019 (before the 2019 Yamagata-Oki earthquake). The E–W contraction rate is shown by the colour scale. The bars denote principal contraction strain rate axes. The star indicates the location of the Yamagata-Oki earthquake.



Fig. 10. Distance change in the E–W direction between the two GPS stations (1162 and 0231; blue crosses in Fig. S1) near the source region of the 2019 Yamagata-Oki earthquake from Jan. 1, 2007 to Sep. 30, 2019. The positive sign indicates the increase in the distance between the two stations. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

the E–W extension strain rate spread over northeast Japan. Near the source region of the Yamagata-Oki earthquake, this E–W extension strain rate progressed until the occurrence of the mainshock due to the postseismic deformation.

The observed upward hypocenter migrations, together with the earthquake occurrence in the stress shadow and in the E–W extension strain rate field, suggest that the Yamagata-Oki earthquake was caused by a reduction in the fault strength due to rising crustal fluids from the deep part. Moreover, if the deep extension of the fault actually existed in the lower crust, fluids may have promoted aseismic deformations in that deep part of the fault. Such localized aseismic deformations beneath the fault may have partly contributed to the occurrence of this earthquake and the formation of the active deformation zone along the eastern margin of the Japan Sea.

In this study, we attempted to constrain the occurrence mechanism of the 2019 Yamagata-Oki earthquake by considering the occurrence of an earthquake in the stress shadow, on the basis that some local effects are required to cause such an earthquake. Fluid migrations and other aseismic processes likely contribute to the occurrence of earthquakes in more general situations.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgement

We deeply thank the editor (Kelin Wang), and three anonymous reviewers for their constructive comments, which helped to improve the manuscript. This study used hypocenters and P-and S-wave arrival time data reported in the unified catalogue of the JMA (https://www.data.jma.go.jp/svd/eqev/data/bulletin/index_e.html). The seismograms were collected and stored by JMA, national universities, and National Research Institute for Earth Science and Disaster Resilience (http://www.hinet.bosai.go.jp/?LANG = en). The figures in this paper were created using GMT (Wessel and Smith, 1998). This research was supported by JSPS KAKENHI Grant Number JP 17K1437 and 17H05309.

Appendix A. Supplementary data

Supplementary data to this article can be found online at https://

doi.org/10.1016/j.tecto.2020.228609.

References

- Asano, Y., Saito, T., Ito, Y., Shiomi, K., Hirose, H., 2011. Spatial distribution and focal mechanisms of aftershocks of the 2011 off the Pacific coast of Tohoku Earthquake. Earth Planets Sp. 63, 669–673. https://doi.org/10.5047/eps.2011.06.016.
- Bird, P., 2003. An updated digital model of plate boundaries. Geochem. Geophys. Geosyst. 4. https://doi.org/10.1029/2001GC000252.
- Calais, E., Camelbeeck, T., Stein, S., Liu, M., Craig, T.J., 2016. A new paradigm for large earthquakes in stable continental plate interiors. Geophys. Res. Lett. 43, 10–621.
- Dahm, T., 1996. Relative moment tensor inversion based on ray theory: theory and synthetic tests. Geophys. J. Int. 124, 245–257. https://doi.org/10.1111/j.1365-246X 1996.tb06368.x.
- Das, S., Henry, C., 2003. Spatial relation between main earthquake slip and its aftershock distribution. Rev. Geophys. 41. https://doi.org/10.1029/2002RG000119.
- Ebel, J.E., Chambers, D.W., 2016. Using the locations of $M \ge 4$ earthquakes to delineate the extents of the ruptures of past major earthquakes. Geophys. J. Int. 207, 862–875. https://doi.org/10.1093/gji/ggw312.
- Eshelby, J.D., 1957. The determination of the elastic field of an ellipsoidal inclusion, and related problems. Proc. R. Soc. Lond. A. Math. Phys. Sci. 376–396. https://doi.org/ 10.1098/rspa.1983.0054.
- Frank, W.B., Poli, P., Perfettini, H., 2017. Mapping the rheology of the central Chile subduction zone with aftershocks. Geophys. Res. Lett. 44, 5374–5382.
- Fukuyama, E., Ishida, M., Dreger, D.S., Kawai, H., 1998. Automated seismic moment tensor determination by using on-line broadband seismic waveforms. Jishin 51, 149–156.
- Gibbons, S.J., Ringdal, F., 2006. The detection of low magnitude seismic events using array-based waveform correlation. Geophys. J. Int. 165, 149–166. https://doi.org/ 10.1111/j.1365-246X.2006.02865.x.
- Hanks, T.C., Kanamori, H., 1979. A moment magnitude scale. in:. J. Geophys. Res. B Solid Earth 2348–2350. https://doi.org/10.1029/JB084iB05p02348.
- Hasegawa, A., Nakajima, J., 2004. Geophysical constraints on slab subduction and arc magmatism. In: Sparks, R.S.J., Hawkesworth, C.J. (Eds.), Geophysical Monograph Series. AGU, pp. 81–93. https://doi.org/10.1029/150GM08.
- Hasegawa, A., Umino, N., Takagi, A., 1978. Double-planed structure of the deep seismic zone in the northeastern Japan arc. Tectonophysics 47, 43–58. https://doi.org/10. 1016/0040-1951(78)90150-6.
- Hasegawa, A., Nakajima, J., Umino, N., Miura, S., 2005. Deep structure of the northeastern Japan arc and its implications for crustal deformation and shallow seismic activity. Tectonophysics 403, 59–75. https://doi.org/10.1016/j.tecto.2005.03.018.
- Heki, K., Miyazaki, S., Takahashi, H., Kasahara, M., Kimata, F., Miura, S., Vasilenko, N.F., Ivashchenko, A., An, K.-D., 1999. The Amurian Plate motion and current plate kinematics in eastern Asia. J. Geophys. Res. Solid Earth 104, 29147–29155. https:// doi.org/10.1029/1999JB900295.
- Helmstetter, A., Sornette, D., 2002. Diffusion of epicenters of earthquake aftershocks, Omori's law, and generalized continuous-time random walk models. Phys. Rev. E Stat. Phys. Plasmas Fluids Relat. Interdiscip. Top. 66, 24. https://doi.org/10.1103/ PhysRevE.66.061104.
- Hsu, Y.-J., Simons, M., Avouac, J.-P., Galetzka, J., Sieh, K., Chlieh, M., Natawidjaja, D., Prawirodirdjo, L., Bock, Y., 2006. Frictional afterslip following the 2005 Nias-Simeulue earthquake. Sumatra. Sci. 312, 1921–1926.
- Hu, Y., Bürgmann, R., Uchida, N., Banerjee, P., Freymueller, J.T., 2016. Stress-driven relaxation of heterogeneous upper mantle and time-dependent afterslip following the 2011 Tohoku earthquake. J. Geophys. Res. Solid Earth 121, 385–411. https://doi. org/10.1002/2015JB012508.
- Iinuma, T., Hino, R., Kido, M., Inazu, D., Osada, Y., Ito, Y., Ohzono, M., Tsushima, H., Suzuki, S., Fujimoto, H., Miura, S., 2012. Coseismic slip distribution of the 2011 off the Pacific Coast of Tohoku Earthquake (M9.0) refined by means of seafloor geodetic data. J. Geophys. Res. Solid Earth 117. https://doi.org/10.1029/2012JB009186.
- Iio, Y., Kobayashi, Y., 2002. A physical understanding of large intraplate earthquakes. Earth Planets Sp. 54, 1001–1004. https://doi.org/10.1186/BF03353292.
- Kagan, Y.Y., 1991. 3-D rotation of double-couple earthquake sources. Geophys. J. Int. 106, 709–716. https://doi.org/10.1111/j.1365-246X.1991.tb06343.x.
- Kato, N., 2004. Interaction of slip on asperities: Numerical simulation of seismic cycles on a two-dimensional planar fault with nonuniform frictional property. J. Geophys. Res. Solid Earth 109, 1–17. https://doi.org/10.1029/2004JB003001.
- Kato, N., 2007. Expansion of aftershock areas caused by propagating post-seismic sliding. Geophys. J. Int. 168, 797–808. https://doi.org/10.1111/j.1365-246X.2006.03255.x.
 Kobavashi, Y., 1983. The start of "subduction" of plate. Earth Monthly 3, 510–518.
- Kosugain, F., Poor Inc start of Automatical of plate functional plate function of the order Kosuga, M., 2014. Seismic activity near the Moriyoshi-zan volcano in Akita Prefecture, northeastern Japan: implications for geofluid migration and a midcrustal geofluid reservoir. Earth, Planets Sp 66. https://doi.org/10.1186/1880-5981-66-77.
- Lay, T., Ammon, C.J., Hutko, A.R., Kanamori, H., 2010. Effects of kinematic constraints on teleseismic finite-source rupture inversions: Great Peruvian earthquakes of 23 June 2001 and 15 August 2007. Bull. Seismol. Soc. Am. 100, 969–994.
- Liu, L., Zoback, M.D., 1997. Lithospheric strength and intraplate. Tectonics 16 (4), 585–595.
- Matsumoto, Y., Yoshida, K., Hasegawa, A., Matsuzawa, T., 2020. Fault-valve behavior estimated from intensive foreshock and aftershock activity in the 2017 M 5.3 Kagoshima Bay, Kyushu, southern Japan, earthquake sequence. Earth Sp. Sci. Open Arch. 30. https://doi.org/10.1002/essoar.10503222.1.
- Mendoza, C., Hartzell, S.H., 1988. Aftershock patterns and main shock faulting. Bull. Seismol. Soc. Am. 78, 1438–1449.
- Meneses-Gutierrez, A., Sagiya, T., 2016. Persistent inelastic deformation in central Japan

revealed by GPS observation before and after the Tohoku-oki earthquake. Earth Planet. Sci. Lett. 450, 366–371. https://doi.org/10.1016/j.epsl.2016.06.055.

- Nakamura, K., 1983. Possible nascent trench along the Eastern Japan sea as the convergent boundary between Eurasian and North American plates. Bull. Earthq. Res. Inst., Univ. Tokyo 58, 711–722.
- National Research Institute for Earth Science and Disaster Resilience, 2019a. NIED Hi-net. Natl. Res. Inst. Earth Sci. Disaster Resil. https://doi.org/10.17598/NIED.0003.
- National Research Institute for Earth Science and Disaster Resilience, 2019b. NIED K-NET, Kik-net. Natl. Res. Inst. Earth Sci. Disaster Resil. https://doi.org/10.17598/ NIED.0004.
- Nur, A., Booker, J.R., 1972. Aftershocks caused by pore fluid flow? Science 175, 885–887. https://doi.org/10.1126/science.175.4024.885.
- Ohtake, M., 1995. A seismic gap in the eastern margin of the Sea of Japan as inferred from the time-space distribution of past seismicity. Island Arc 4, 156–165.
- Okada, T., Matsuzawa, T., Umino, N., Yoshida, K., Hasegawa, A., Takahashi, H., Yamada, T., Kosuga, M., Takeda, T., Kato, A., Igarashi, T., Obara, K., Sakai, S., Saiga, A., Iidaka, T., Iwasaki, T., Hirata, N., Tsumura, N., Yamanaka, Y., Terakawa, T., Nakamichi, H., Okuda, T., Horikawa, S., Katao, H., Miura, T., Kubo, A., Matsushima, T., Goto, K., Miyamachi, H., 2015. Hypocenter migration and crustal seismic velocity distribution observed for the inland earthquake swarms induced by the 2011 Tohoku-Oki earthquake in NE Japan: implications for crustal fluid distribution and crustal permeability. Geofluids 15, 293–309. https://doi.org/10.1111/gfl.12112.
- Okamura, Y., Watanabe, M., Morijiri, R., Satoh, M., 1995. Rifting and basin inversion in the eastern margin of the Japan Sea. Island Arc 4, 166–181. https://doi.org/10.1111/ j.1440-1738.1995.tb00141.x.
- Ozawa, S., Nishimura, T., Suito, H., Kobayashi, T., Tobita, M., Imakiire, T., 2011. Coseismic and postseismic slip of the 2011 magnitude-9 Tohoku-Oki earthquake. Nature 475, 373–376. https://doi.org/10.1038/nature10227.
- Parotidis, M., Shapiro, S.a., Rothert, E., 2005. Evidence for triggering of the Vogtland swarms 2000 by pore pressure diffusion. J. Geophys. Res. Solid Earth 110, 1–12. https://doi.org/10.1029/2004JB003267.
- Peng, Z., Zhao, P., 2009. Migration of early aftershocks following the 2004 Parkfield earthquake. Nat. Geosci. 2, 877–881. https://doi.org/10.1038/ngeo697.
- Perfettini, H., Frank, W.B., Marsan, D., Bouchon, M., 2018. A model of aftershock migration driven by afterslip. Geophys. Res. Lett. 45, 2283–2293. https://doi.org/10. 1002/2017GL076287.
- Rice, J.R., 1992. Fault stress states, pore pressure distributions, and the weakness of the San Andreas fault. In: Evans, B., Wong, T.- f. (Eds.), Fault Mechanics and Transport Properties of Rocks, Int. Geophys. Set. 51. Academic, San Diego, Calif., pp. 9885–9907.
- Sagiya, T., Miyazaki, S., Tada, T., 2000. Continuous GPS array and present-day crustal deformation of Japan. Pure Appl. Geophys. 157, 2303–2322.
- Schaff, D.P., Waldhauser, F., 2010. One magnitude unit reduction in detection threshold by cross correlation applied to parkfield (California) and China seismicity. Bull. Seismol. Soc. Am. 100, 3224–3238. https://doi.org/10.1785/0120100042.
- Shapiro, S.A., Huenges, E., Borm, G., 1997. Estimating the crust permeability from fluidinjection-induced seismic emission at the KTB site. Geophys. J. Int. 131, F15–F18. https://doi.org/10.1111/j.1365-246X.1997.tb01215.x.
- Shelly, D.R., Moran, S.C., Thelen, W.A., 2013. Evidence for fluid-triggered slip in the 2009 Mount Rainier, Washington earthquake swarm. Geophys. Res. Lett. 40, 1506–1512. https://doi.org/10.1002/grl.50354.
- Shen, Z.-K., Jackson, D.D., Ge, B.X., 1996. Crustal deformation across and beyond the Los Angeles basin from geodetic measurements. J. Geophys. Res. Solid Earth 101, 27957. https://doi.org/10.1029/96JB02544.
- Sibson, R., 1992. Implications of fault-valve behaviour for rupture nucleation and recurrence. Tectonophysics 211, 283–293. https://doi.org/10.1016/0040-1951(92) 90065-E.
- Sun, T., Wang, K., Iinuma, T., Hino, R., He, J., Fujimoto, H., Kido, M., Osada, Y., Miura, S., Ohta, Y., Others, 2014. Prevalence of viscoelastic relaxation after the 2011 Tohokuoki earthquake. Nature 514, 84–87.
- Terakawa, T., Hashimoto, C., Matsu'ura, M., 2013. Changes in seismic activity following the 2011 Tohoku-oki earthquake: effects of pore fluid pressure. Earth Planet. Sci. Lett. 365, 17–24. https://doi.org/10.1016/j.epsl.2013.01.017.
- Tomita, F., Iinuma, T., Ohta, Y., Hino, R., Kido, M., Uchida, N., 2020. Improvement on spatial resolution of a coseismic slip distribution using postseismic geodetic data

through a viscoelastic inversion. Earth Planets Sp. 72, 1-24.

- Waldhauser, F., Ellsworth, W.L., 2000. A double-difference earthquake location algorithm: method and application to the Northern Hayward fault, California. Bull. Seismol. Soc. Am. 1353–1368.
- Wei, D., Seno, T., 1998. Determination of the Amurian plate motion. Mantle Dyn. Plate Interact. East Asia 27, 337–346.
- Wells, D.L., Coppersmith, K.J., 1994. New empirical relationships among magnitude, rupture length, rupture width, rupture area, and surface displacement. Bull. Seismol. Soc. Am. 84, 974–1002.
- Wessel, P., Smith, W.H.F., 1998. New, improved version of generic mapping tools released. EOS Trans. Am. Geophys. Union 79, 579. https://doi.org/10.1029/ 98E/000426.
- Wesson, R.L., 1987. Modelling aftershock migration and afterslip of the San Juan Bautista, California, earthquake of October 3, 1972. Tectonophysics 144, 215–229. https://doi.org/10.1016/0040-1951(87)90019-9.
- Wetzler, N., Lay, T., Brodsky, E.E., Kanamori, H., 2018. Systematic deficiency of aftershocks in areas of high coseismic slip for large subduction zone earthquakes. Sci. Adv. 4. https://doi.org/10.1126/sciadv.aao3225.
- Woessner, J., Wiemer, S., 2005. Assessing the quality of earthquake catalogues: estimating the magnitude of completeness and its uncertainty. Bull. Seismol. Soc. Am. 95, 684–698.
- Woessner, J., Schorlemmer, D., Wiemer, S., Mai, P.M., 2006. Spatial correlation of aftershock locations and on-fault main shock properties. J. Geophys. Res. Solid Earth 111. https://doi.org/10.1029/2005JB003961.
- Yamashita, T., Knopoff, L., 1987. Models of aftershock occurrence. Geophys. J. R. Astron. Soc. 91, 13–26. https://doi.org/10.1111/j.1365-246X.1987.tb05210.x.
- Yoshida, K., Hasegawa, A., 2018a. Sendai-Okura earthquake swarm induced by the 2011 Tohoku-Oki earthquake in the stress shadow of NE Japan: detailed fault structure and hypocenter migration. Tectonophysics 733, 132–147. https://doi.org/10.1016/j. tecto.2017.12.031.
- Yoshida, K., Hasegawa, A., 2018b. Hypocenter migration and seismicity pattern change in the Yamagata-Fukushima border, NE Japan, caused by fluid movement and pore pressure variation. J. Geophys. Res. Solid Earth 123, 5000–5017. https://doi.org/10. 1029/2018JB015468.
- Yoshida, K., Hasegawa, A., Okada, T., Iinuma, T., Ito, Y., Asano, Y., 2012. Stress before and after the 2011 great Tohoku-oki earthquake and induced earthquakes in inland areas of Eastern Japan, Geophys. Res. Lett. https://doi.org/10.1029/2011gl049729.
- Yoshida, K., Hasegawa, A., Okada, T., Iinuma, T., 2014. Changes in the stress field after the 2008 M7.2 Iwate-Miyagi Nairiku earthquake in Northeastern Japan. J. Geophys. Res. Solid Earth 119, 9016–9030. https://doi.org/10.1002/2014JB011291.
- Yoshida, K., Hasegawa, A., Okada, T., 2015. Spatial variation of stress orientations in NE Japan revealed by dense seismic observations. Tectonophysics 647, 63–72. https:// doi.org/10.1016/j.tecto.2015.02.013.
- Yoshida, K., Hasegawa, A., Yoshida, T., 2016. Temporal variation of frictional strength in an earthquake swarm in NE Japan caused by fluid migration. J. Geophys. Res. Solid Earth 121, 5953–5965. https://doi.org/10.1002/2016JB013022.
- Yoshida, K., Hasegawa, A., Yoshida, T., Matsuzawa, T., 2019a. Heterogeneities in stress and strength in Tohoku and its relationship with earthquake sequences triggered by the 2011 M9 Tohoku-Oki earthquake. Pure Appl. Geophys. https://doi.org/10.1007/ s00024-018-2073-9.
- Yoshida, K., Saito, T., Emoto, K., Urata, Y., Sato, D., 2019b. Rupture directivity, stress drop, and hypocenter migration of small earthquakes in the Yamagata-Fukushima border swarm triggered by upward pore-pressure migration after the 2011 Tohoku-Oki earthquake. Tectonophysics 769. https://doi.org/10.1016/j.tecto.2019.228184.
- Yoshida, K., Taira, T., Matsumoto, Y., Saito, T., Emoto, K., Matsuzawa, T., 2020. Stress release process along an intraplate fault analogous to the plate boundary: a case study of the 2017 M5.2 Akita-Daisen earthquake, NE Japan. J. Geophys. Res. Solid Earth 125. https://doi.org/10.1029/2020JB019527.
- Yukutake, Y., Honda, R., Harada, M., Tanada, T., Yoshida, A., 2011. Fluid-induced swarm earthquake sequence revealed by precisely determined hypocenters and focal mechanisms in the 2009 activity at Hakone volcano, Japan. J. Geophys. Res. Earth 116, 13. https://doi.org/10.1029/2010jb008036.
- Zhao, D., Huang, Z., Umino, N., Hasegawa, A., Yoshida, T., 2011. Seismic imaging of the Amur-Okhotsk plate boundary zone in the Japan Sea. Phys. Earth Planet. Inter. 188, 82–95. https://doi.org/10.1016/j.pepi.2011.06.013.